



Analysis of Requisite Quality of Cassava and Temperature for Baking Cookies

SARIN NEANG*

*Royal University of Agriculture, Phnom Penh, Cambodia
Email: neangsarin@gmail.com*

VITOU ORNG

Royal University of Agriculture, Phnom Penh, Cambodia

VOUCHSIM KONG

Royal University of Agriculture, Phnom Penh, Cambodia

SAROM MEN

Royal University of Agriculture, Phnom Penh, Cambodia

LINDA FORRESTER

Education Advisor, Australia

BORARIN BUNTONG

Royal University of Agriculture, Phnom Penh, Cambodia

Received 16 December 2012 Accepted 10 June 2013 (*Corresponding Author)

Abstract Cookies are a kind of food product made from flour and other ingredients such as butter, egg, sugar, milk, and baking soda. This research was conducted to find the optimum temperature and the optimum proportion of cassava flour required in cookie production. The experiment was divided into nine treatments with three replications, different temperatures included 160 °C, 170 °C ; 180 °C and proportions of cassava flour trialed were 50 %, 75 %; 100 %. The processing and sensory testing were conducted in the food processing laboratory of the Faculty of Agro-Industry, Royal University of Agriculture (RUA) while the analysis of chemicals was done in the Industrial Laboratory Centre of Cambodia (ILCC). The results show that using 50%, 70% and 100% 5 % and 100 % of cassava flour at the temperature of 160 °C, 170 °C and 180 °C affected the chemical properties of cookie products such as ash, moisture, protein, fat, sugar, carbohydrate, and energy. Also, they affected the color, texture, odor, as well as flavor and crispness, factors which all related to consumers identifying cookies as their favorites. Among nine treatments, the second treatment which used 50% of cassava flour at the temperature of 170 °C was the best treatment; whereas, the ninth treatment which used 100% of cassava flour at the temperature of 180 °C had the lowest quality. Therefore, results show that 50 % of cassava flour baked at the temperature of 170 °C for 8 minutes is the optimum configuration to use in cookie production.

Keywords processing, temperature, sensory test, cassava, cookies

INTRODUCTION

Cassava is a kind of plant that mostly grows in tropical area in South America. The root of Cassava has poisonous substances, but the rest of the plant is edible and provides a lot of essential nutrients. Cassava is rich in Carbohydrate and low lipids that play a role in providing energy to the human body. Energy from Cassava is related to the density of calories, where 100 gram provides 160 calories (Willett, 2005). Cassava is known and grown by Cambodian farmers since 19th century. It was grown for starch production for commercial purpose, livestock feed and used for human.

Before 1970, this crop was planted in small areas, especially on black soil, along the river, and upland areas and red soil in some areas of the country.

Between 1975 and 1979, this crop becomes a staple, and people eat cassava instead of rice. This is a very popular crop among farmers in southeastern and eastern areas of the country, especially for people living in upland areas. However, some farmers living in lowland areas also grow this crop on the space around their house, and they consume it as their food (MAFF, 2003).

The cassava yield increased from 535,600 tons in 2005 to 2 million tons in 2007. This is the result of increasing lands for growing cassava crops. Cassava was planted on 30,000 hectares in 2005 and 1,000,000 hectares in 2007 and the yield rose from 17.9 tons/ha to 20.5 tons/ha (CIAT, 2008). Cassava has an important role in the development of rural areas because it is the raw material in industrial productions such as animal feed, starch, ethanol, bioethanol and other food processes in bakery production, such as ice cream, cakes, and biscuits (Willett, 2005).

OBJECTIVE

This research was aimed at finding the optimum temperature and the amount of cassava flour required in cookie production.

METHODOLOGY

Study site

This research included processing cookies, analyzing chemical contents and sensory testing and was conducted at the laboratory of the faculty of Agro-Industry, Royal University of Agriculture (RUA). The analysis of chemical contents in processed cookies was done at the Industrial Laboratory Centre of Cambodia (ILCC), Phnom Penh, while the sensory test was done at RUA. The research study was conducted between May and August, 2012.

Experimental design and treatments

The factorial Complete Randomized Design (CRD) was used in this experimental method; there are three factors including temperature (160°C, 170 °C and 180 °C), cassava flour (50%, 75% and 100%), and wheat flour (50% and 25%) as shown in table 1.

Table 1 Divisions of experimental treatments

Treatments	Temperature (°C)	Cassava flour (%)	Wheat flour (%)
T ₁	160		
T ₂	170	50	50
T ₃	180		
T ₄	160		
T ₅	170	75	25
T ₆	180		
T ₇	160		
T ₈	170	100	00
T ₉	180		

Raw materials of the cookies were cassava flour, wheat flour, sugar, egg, butter, baking soda, blue berries and milk. 11% of egg white was mixed by a hand mixer and then 31. 25% of butter was added in the egg port. 31. 25% of sugar, 6.25% of milk, 16% of blue berries and 1% of baking soda were subsequently mixed in the egg and butter port. The cassava flour and wheat flour were added and compressed into cookie dough 1 cm thick before baking in the oven for eight minutes in each temperature that had already been set. Noticeably, before using the oven, it needed to be