



Dietary Exposure Assessment of Benzylpenicillin Residue in Pork Consumed by Age and Gender Groups in the Philippines

VENN VUTEY*

*Faculty of Veterinary Medicine, Royal University of Agriculture, Cambodia
Email: vennvutey@gmail.com*

LOINDA R. BALDIAS

College of Veterinary Medicine, University of the Philippines Los Baños, Philippines

BILLY P. DIVINA

College of Veterinary Medicine, University of the Philippines Los Baños, Philippines

RIO JOHN T. DUCUSIN

College of Veterinary Medicine, University of the Philippines Los Baños, Philippines

Received 16 December 2012 Accepted 30 January 2013 (*Corresponding Author)

Abstract Exposure assessment is one of the most important key components of the risk assessment process. Dietary intake of significant amounts of residue can lead to adverse health effects and development of antimicrobial resistance in the population. The study was conducted to determine exposure risk to antibiotic drug residues in pork consumed in the Philippines. The specific aim was to estimate dietary exposure of Benzylpenicillin residues by age and gender groups. Parameters such as food consumption, substances residue, body weight, ages, and gender groups were gathered from local and international institutions. Mathematical equations were used to calculate for Dietary Exposure from amount of drug residue multiplied by the mean food consumption and adjusted by body weight. The present study; dietary exposure of Benzylpenicillin residue for children from 1 to <3 years old had the highest exposed followed by adolescents, adults and infants while males were significantly higher than females ($p < 0.05$). This is the first attempt to determine risk assessment of dietary exposure to antibiotic residue of different population groups in the Philippines. Improvement of mathematical models used in this study is proposed to better prioritize exposure assessment models for veterinary drug residues to ensure the safety of food produced from farm to table.

Keywords exposure assessment, drug residues, age-gender groups, consumption

INTRODUCTION

The use of risk assessment has gained steadily in importance and recognition as a scientifically-based approach for the development of food safety and quality standards. During recent years there has been increasing use of the word “risk” in connection with food safety (Sumner et al., 2004). Consequently, a risk assessment of the food safety implications (Gehring et al., 2005) may be made on the potential harm or risks of veterinary drug residues to the human consumer. The importance of risk assessment lies not only in its ability to estimate human risk, but also in its use as a framework for organizing data, as well as for allocating responsibility for analysis. As such according to World Health Organization (1995), it is important to understand that risk assessment is a process that can include a variety of mathematical models to estimate risks. Risk assessments also promote international trade by addressing new safety and quality challenges (Graf, 2005).

Exposure assessment is among the components of risk assessment (Codex Alimentarius Commission, 1999). Dietary exposure to veterinary drugs occurs when residues remain in food

products derived from animals treated with a veterinary drug (Health Canada, 2003a). Dietary exposure models have been developed to predict human exposures to chemicals resulting from the consumption of contaminated food and residue levels and also for substances naturally occurring in them (Fryer, 2006). In 2003b, Health Canada primer on “Assessing exposure from pesticides in food” indicated that risk exposure from food can either result in short-term (i.e., acute) or long term (i.e., chronic) effects. In actual risk equations, it is possible that toxicity may be computed utilizing the estimation of consumer intake of drug residues in foods of animal origin, hence, this study on exposure assessment.

Thorough literature search revealed no available study on exposure assessment on veterinary drug residues in the Philippines. With the increase in reports of occurrence of risk of veterinary drug residues in foods of animal origin in the last two decades (Health Canada, 2003a; Sumner et al., 2004), there is a strong need to study exposure assessment of antibiotic residues. To date, there is no local study conducted on the exposure risk assessment for residual antibiotics in the Philippines. Thus, this study aims to make a “Dietary Exposure Assessment of Benzylpenicillin Residues in Pork Consumed by Age and Gender Groups in the Philippines”.

The overall objective of the study is to determine exposure risk to antibiotic drug residues in pork consumed in the Philippines. The specific aim is to estimate the dietary exposure of Benzylpenicillin residues by ages and gender groups.

METHODOLOGY

The study was conducted using secondary data collected from local and international institutions for estimating dietary exposure assessment. For instance, local data like the report on veterinary drug residue (2003-2008) from National Meat Inspection Service, recommendation on Maximum Residue Limit (MRL) published by Bureau of Aquaculture Fisheries and Product Standard, the report of Food Consumption Survey in 2003 and Recommended Energy and Nutrient Intake in 2002 published by Food and Nutrition Research Institute (FNRI, 2003), the development of antimicrobial resistance from Research Institute for Tropical Medicine, Philippines. International data came from Codex Alimentarius Commission (CAC) on amount of antibiotic drug residues, the procedural guidelines on residues of veterinary drugs in food from JECFA, and from an updated report of the 32nd session of the Codex Veterinary Drug Residues in Food 2009.

Dietary modelling combines food consumption data with food chemical concentration data to estimate dietary exposure to food chemicals, or intake of nutrients. The Eq. (1) below, the basic dietary exposure model is of the form:

$$\text{Dietary Exposure} = \frac{\text{Food Consumption} \times \text{Amount of drug residue}}{\text{Body weight}} \quad (1)$$

where food consumption is the amount of food people eating per capita per day expressed in mg/kg/day and amount of drug residue expressed in µg/kg BW/day.

RESULTS

The benzylpenicillin antibiotic have been test for consecutive years from 2006 to 2008 in the Philippines. The number of violations was determined using Microbial Inhibition Test and was measured using caliper method (NMIS, 2008). Positive benzylpenicillin residue results in NMIS were assumed to equivalent to HPLC LOQ 400 µg/kg (Heitzman, 1995) and dietary exposure was calculated using the following equation (3) which regarded computed using equation (2).

Eq. (2) the Mean Food Consumption Difference Age Groups is as follows;

$$\text{The mean food consumption difference age groups} = \frac{\text{average weight difference age groups} \times \text{the mean food consumption per capita in the Philippines}}{\text{average weights}} \quad (2)$$

Eq. 3 Dietary Exposure of Benzylpenicillin Residue is as follows;

$$DE_{PEN} = \frac{\text{amount of benzylpenicillin residues of 3\% positive results} \times \text{mean food consumption difference age groups}}{\text{average weights differences age groups}} \quad (3)$$

Based on an update as of the 32nd session of the Codex Alimentarius Commission in 2009 first draft prepared by Dr. Macneil, a typical recommended dose by intramuscular injection (IM) in swine, cattle, horses, and sheep of a 300,000 unit/mL formulation is 6,600 units/mL BW. As a feed additive, a typical dosage for swine or poultry is 55 mg/kg in the diet. Intramammary treatment is typically by administration of 100,000 units per quarter. In the studies reported, 1 mg procaine benzylpenicillin is equivalent to 1667 IU (international units).

As at 50th report of JECFA in 1999, the Committee considered residues of procaine benzylpenicillin to be equivalent to residues of benzylpenicillin. Dietary Exposure (DE) was calculated using equation (1) following the mean one-day per capita food consumption in the Philippines and by island groups, food groups and adjusted 55-kg person. The DE of Benzylpenicillin Residue was calculated by the amount of benzylpenicillin of 3% positive results multiplied the food consumption difference age groups and adjusted by average weight difference age groups.

Table 1 shows that DE of benzylpenicillin residue for children from 1 to 3 years old had the highest value at 0.002923 µg/kg bw/day. Such value was calculated as 5 µg/kg of amount residues × mean one-day per capita pork consumption of 0.0076 kg ÷ 13 kg of average children weights. Likewise, average DE for children (0.002913 µg/kg bw/day) was higher than adolescents (0.0029096 µg/kg bw/day), followed by adults (0.002909 µg/kg bw/day) and infants (0.002903 µg/kg bw/day). Also, DE for males was significantly lower than females (p-value=0.034).

Table 1 Dietary exposure of benzylpenicillin residue of 3% of positive results intake of fresh pork in the Philippines

Age Groups	Average Weight (kg)	Amount Residue (µg/kg)	Mean One-Day Per Capita Pork Consumption (g)	Dietary Exposure (µg /kg bw/day)
1. Infants, month				
Birth- <6 (3)	6	5	3.5	0.002917
6-<12 (9)	9		5.2	0.002889
2. Children, year				
1-3 (2.5)	13		7.6	0.002923
4-6 (5.5)	19	5	11.1	0.002921
7-9 (8.5)	24		13.9	0.002896
3. Adolescents, M, year				
10-12 (11.5)	34		19.8	0.002911
13-15 (14.5)	50	5	29.1	0.002910
16-18 (17.5)	58		33.7	0.002905
4. Adolescents, F, year				
10-12 (11.5)	35		20.4	0.002914
13-15 (14.5)	49	5	28.5	0.002908
16-18 (17.5)	50		29.1	0.002910
5. Adults, M, year				
19-49	59		34.3	0.002907
50-64	59	5	34.3	0.002907
65 and over	59		34.3	0.002907
6. Adults, F, year				
19-49	51		29.7	0.002911
50-64	51	5	29.7	0.002911
65 and over	51		29.7	0.002911