



Estimated Maximum Daily Intake of Streptomycin Residue in Pork Consumed by Age and Gender Groups in the Philippines

VENN VUTEY*

Faculty of Animal Science and Veterinary Medicine, Royal University of Agriculture, Phnom Penh, Cambodia

Email: vennvutey@gmail.com

LOINDA R. BALDIAS

CVM-UPLB, Philippines

BILLY P. DIVINA

CVM-UPLB, Philippines

RIO JOHN T. DUCUSIN

Chair of Department of Veterinary Clinical Sciences, CVM-UPLB, Philippines

Received 14 December 2011 Accepted 7 February 2012 (*: Corresponding Author)

Abstract Dietary intake of significant amounts of residues can lead to adverse health effects and the development of antimicrobial resistance in the population. This study was conducted to determine dietary intake of antibiotic drug residues in pork consumed in the Philippines. The specific aim was to estimate maximum daily intake of streptomycin residue ($EMDI_{STC}$) by age and gender groups. Parameters such as maximum residue limits (MRL), 90th percentile food consumption, body weight, age and gender groups were gathered from local and international institutions. A mathematical equation was used to calculate $EMDI_{STC}$ from MRL multiplied by the 90th percentile food consumption and adjusted by body weight. In the present study, the $EMDI_{STC}$ for infants from birth to less than 12 months of age had the highest intake ratio of streptomycin residue followed by children, adolescents and adults where males were significantly higher than females ($p < 0.05$). Based on the findings of the present study, it is concluded that streptomycin detected in pork affects infants from birth to less than 12 months that are more likely to consume it and more vulnerable due to physical activity. This is the first attempt to estimate dietary intake of antibiotic residues in the Philippines. Improvement of mathematical models used in this study is proposed to prioritize better models for veterinary drug residues to ensure the safety of food produced from farm to table.

Keywords: dietary intake, pork, antibiotic residue, age-gender group, Philippines

INTRODUCTION

Food needs to be consumed for maintenance of human functions. Food has to be safe for human consumption. Food safety is imperative for the development and maintenance of a healthy population (Titus, 2007). Consequently, health effects and safety aspects of food are important issues for today's consumers. There are increasing public health concerns that drug residues and their metabolites could be found in meat and other foods of animal origin may and cause adverse effects on consumers' health (EFSA, 2007).

Through time livestock production has evolved from small scale to large scale integrative farming (NMIS, 2006). This required the increased use of veterinary drugs. These are used to cure or prevent diseases in animals, to increase feed efficiency or growth rate, and to sedate animals in order to minimize the effect of stress (Botsoglou and Flatouris, 2001; Doyle, 2006). With the

widespread use of veterinary drugs in animal production, there is global concern about the consumption of foods of animal origin that may contain residues and their possible adverse effects on human health (Huss et al., 2004). For instance, with the use of antimicrobials in animal production, there is a possibility for the creation of antibiotic-resistant pathogens in animals that may be transferred to man (FSIS, 2000; Doyle, 2006). Boden (2005) cited that drug residues in food are regarded as very important from the public health's point of view. Moreover, there has been concern about carryover of veterinary drug residues in meat, eggs, and milk to people consuming these foods (Botsoglou and Flatouris, 2001; Doyle, 2006). Potential hazards associated with the presence of veterinary drug residues in edible tissues have been reported to cause toxic or allergic reactions, anaphylactic reactions, headache, and severe aplastic anaemia (WHO, 1999; Kelly, 2005).

Thorough literature search revealed no available study on estimated antibiotic drug residues in pork consumed using mathematical equations in the Philippines. With the increase in reports of occurrence of risk of veterinary drug residues in foods of animal origin in the last two decades (Health Canada, 2003a, Sumner et al., 2004), there is a strong need to study dietary intake of veterinary drug residues. To date, there is no local study conducted on the prediction for residual antibiotics in the Philippines. Thus, this study aims to derive an estimated maximum daily intake of streptomycin residues in pork consumed by age and gender groups in the Philippines.

The overall objective of the study is to determine dietary intake of antibiotic drug residues in pork consumed in the Philippines. The specific aim is to estimate maximum daily intake of streptomycin residue by age and gender groups.

METHODOLOGY

The study was conducted using secondary data collected from local and international institutions for estimating EMDI (estimated maximum daily intake). For instance, local data like the report on veterinary drug residue (2003-2008) from National Meat Inspection Service (NMIS), recommendation on Maximum Residue Limit (MRL) published by the Bureau of Aquaculture Fisheries and Product Standard (BAFPS), the report of Food Consumption Survey in 2003 and Recommended Energy and Nutrient Intake in 2002 published by Food and Nutrition Research Institute (FNRI), and the development of antimicrobial resistance from Research Institute for Tropical Medicine (RITM). International data came from Codex Alimentarius Commission (CAC) on amount of antibiotic drug residues, the procedural guidelines on residues of veterinary drugs in food from JECFA, and from an updated report of the 32nd session of the Codex Veterinary Drug Residues in Food 2009.

EMDI modelling combines MRL data with 90th percentile food consumption data and body weight factor to estimate EMDI. The basic EMDI model is of the form:

$$\text{EMDI} = [\text{MRL} \times 90^{\text{th}} \text{ percentile food consumption}] \div \text{weight factor}$$

where EMDI is an estimate of drug residue intake expressed in $\mu\text{g}/\text{kg BW}/\text{day}$.

RESULTS

Streptomycin is active against a wide range of gram-negative organisms and some gram-positive pathogens in pigs, cattle and sheep. Streptomycin was evaluated by the Codex Committee at its 12th meeting in 1969, 43rd meeting in November 1994, 48th meeting in 1998, report of the 11th session of the Codex Committee on residues of veterinary drugs in food in September 1998 and updated as of the 32nd session of the Codex Alimentarius Commission in July 2009. The committee proposed a temporary MRL's for streptomycin at 600 $\mu\text{g}/\text{kg}$ (muscle, liver, fat).

The reference weight for adults, 59 kg for males and 51 kg for females, and the average weight in the Philippines, 55 kg (FNRI, 2003) as well as the 90th percentile food consumption per capita in the Philippines and three islands are shown in Table 1.

Table 1 Mean one-day per capita pork consumption (gram, raw, as purchased) 2003 food consumption survey

	Philippines	Luzon	Visayas	Mindanao
Fresh Meat				
Pork				
Mean	31.80	39.89	23.07	18.24
95% CI	31.76, 31.83	39.84, 39.93	23.01, 23.14	18.19, 18.30
P90	108.21	127.31	88.50	57.29

Source: FNRI, 2003

However, the weight difference between age groups: infants, adolescents, adults and older; the 90th percentile food consumption per capita; and average weights in the Philippines were calculated using Eq. (1) below.

$$\text{The 90th percentile food consumption difference age groups} = [\text{average weight difference age groups} \times \text{the 90th percentile food consumption per capita in the Philippines}] \div \text{average weights in the Philippines} \quad (1)$$

Using Eq. (1), the 90th percentile food consumption different age groups was calculated separately by ages (infants, children, adolescents, and adults), gender, average weights in different age groups, and average weight in the Philippines. A summary for computation of the 90th percentile food consumption in the Philippines is shown in Table 2.

Table 2 Summary of 90th percentile pork consumption per capita by age group and average weight in the Philippines, Luzon, Visayas and Mindanao

Age Groups	Weight (kg)	Philippines (g)	Luzon (g)	Visayas (g)	Mindanao (g)
1. Infants, mo					
Birth- <6 (3)	6	12	14	10	6
6-<12 (9)	9	18	21	15	9
2. Children, y					
1-3 (2.5)	1	26	30	21	13
4-6 (5.5)	19	37	44	31	20
7-9 (8.5)	24	47	55	39	25
3. Adolescents, males, y					
10-12 (11.5)	34	67	79	55	35
13-15 (14.5)	50	98	115	81	52
16-18 (17.5)	58	114	134	94	60
4. Adolescents, females, y					
10-12 (11.5)	35	69	81	57	36
13-15 (14.5)	49	96	113	79	51
16-18 (17.5)	50	98	115	81	52
5. Adults, males, y					
19-49	59	116	136	95	61
50-64	59	116	136	95	61
65 and over	59	116	136	95	61
6. Adults, females, y					
19-49	51	100	118	83	53
50-64	51	100	118	83	53
65 and over	51	100	118	83	53

Eq. (2) was used to calculate the EMDI of streptomycin residues by adaptation from the value of Eq. (1) then multiplied by MRL muscle tissue and adjusted by weight factor as recommended by

Codex committee and Philippines National Standard. Then comparison of estimates of streptomycin residues intake was computed.

$$\text{EMDI}_{\text{STC}} = [\text{MRL} \times \text{the } 90^{\text{th}} \text{ percentile food consumption difference age groups}] \div \text{average weights differences age groups} \quad (2)$$

Table 3 summarizes the EMDI of streptomycin residue for all locations. Particulars for the computation of EMDI of streptomycin residue in the Philippines, Luzon, Visayas and Mindanao are shown in Appendix B Tables 2 to 5 (Venn et al., 2010).

Table 3 Summary of EMDI of streptomycin residue intake of fresh pork in the Philippines, Luzon, Visayas and Mindanao

Age Groups	Philippines ($\mu\text{g}/\text{kg}$ BW/day)	Luzon ($\mu\text{g}/\text{kg}$ BW/day)	Visayas ($\mu\text{g}/\text{kg}$ BW/day)	Mindanao ($\mu\text{g}/\text{kg}$ BW/day)
Infants, mo				
Birth- <6 (3)	1.2000	1.4000	1.0000	0.6000
6-<12 (9)	1.2000	1.4000	1.0000	0.6000
Average	1.2000	1.4000	1.0000	0.6000
Children, y				
1-3 (2.5)	1.2000	1.3846	0.9692	0.6000
4-6 (5.5)	1.1684	1.3894	0.9789	0.6315
7-9 (8.5)	1.1750	1.3750	0.9750	0.6250
Average	1.1811	1.3830	0.9743	0.6188
Adolescents, Males, y				
10-12 (11.5)	1.1823	1.3941	0.9705	0.6176
13-15 (14.5)	1.1760	1.3800	0.9720	0.6240
16-18 (17.5)	1.1793	1.3862	0.9724	0.6206
Adolescents, Females, y				
10-12 (11.5)	1.1828	1.3885	0.9771	0.6171
13-15 (14.5)	1.1755	1.3836	0.9673	0.6244
16-18 (17.5)	1.1760	1.3800	0.9720	0.6240
Average	1.1786	1.1854	0.9718	0.6212
Adults, males, y				
19-49				
50-64	1.1796	1.3830	0.9661	0.6203
65 and over	1.1796	1.3830	0.9661	0.6203
	1.1796	1.3830	0.9661	0.6203
Adults, females, y				
19-49	1.1764	1.3882	0.9764	0.6235
50-64	1.1764	1.3882	0.9764	0.6235
65 and over	1.1764	1.3882	0.9764	0.6235
Average	1.1780	1.1856	0.9712	0.6219

Fig. 1 shows that EMDI of streptomycin residues for infants is 1.2 $\mu\text{g}/\text{kg}$ BW/day from birth to less than 12 months of age. 1.2 $\mu\text{g}/\text{kg}$ BW/day was calculated as 600 $\mu\text{g}/\text{kg}$ of MRL multiplied by 90th percentile food consumption infant of 0.012 kg divided by the average weight of infant at 6 kg. It is equivalent to children from one to three years of age. In general, infants had consumed more streptomycin residues than children (1.1811 $\mu\text{g}/\text{kg}$ BW/day) followed by adolescents (1.1786 $\mu\text{g}/\text{kg}$ BW/day) and adults (1.1780 $\mu\text{g}/\text{kg}$ BW/day). Moreover, EMDI for males was significantly higher than that for females ($p < 0.05$).




Fig. 1 Estimated maximum daily intake of streptomycin residue for females and males from different age and weight groups in the Philippines

DISCUSSION

Prediction of EMDI of streptomycin residues from dietary intake of pork using a stepwise approach is a step forward to provide an estimate potential exposure of the population. Hence, analysis of EMDI found that infants from birth to less than 12 months of age had the highest intake of streptomycin residue in the Philippines, followed by children, adolescents and adults.

A similar study on the exposure assessment of chemical from packaging materials in food by Pocas and Hogg (2007) mentioned that food consumption by infants and small children is much higher, based on the body weight, than by conventional adult model of $1 \text{ kg}_{\text{food}}/60 \text{ kg BW}$. In addition, infants and children, because of their higher food consumption rates per kg BW, were generally expected to have a higher dietary daily intake of substances (Kroes et al., 2002). However, the assessment on dietary Melamine exposure, Xu et al. (2009) reported that infants of three months old had the highest intake estimate of melamine and with the increase in age, the intake decreased. Another example was conducted on dietary exposure to copper in the European Union. Steve (2008) reported that in most studies in EU member states, age and sex differences were more apparent than regional differences. Young and middle-aged adults have higher mean intake than elderly or very elderly.

CONCLUSION

Based on findings of the present study, streptomycin residue have been detected in pork which infants from birth to less than 12 months are more exposed to and more at risk, because of their higher food consumption rates per kg of body weight, and they are generally expected to have a higher exposure level due to physical activity.

ACKNOWLEDGMENTS

First, my thanks go to Veterinary Drug Residue Section of NMIS, Quezon City Philippines for allow using the new report of veterinary drug residue monitoring program and laboratory technicians of this study. I am thankful to Antimicrobial Resistance Surveillance Program Team of RITM and National Epidemiology Centre, Muntinlupa City, Philippines. I also wish to thank Food of Nutrition Research Institute Team, who has given the 2003 food consumption survey and percentile data released of fresh pork in the Philippines and RENI-FNRI-DOST, Metro Manila, Philippines.

REFERENCES

- Boden, E. 2005. Black's Veterinary Dictionary, 21st edition.
- Botsoglou, N., and Fletouris, D. 2001. Drug residues in foods. Pharmacology, Food Safety, and Analysis, Aristotle University, Thessaloniki, Greece.
- CAC. 1999. Procedural manual, tenth edition, Joint FAO/WHO Food Standards Programme, Food and Agriculture Organization of the United Nations, 1997. Definitions for "risk management" and "risk communication" were adopted as revised texts at the twenty-third session of the Codex Alimentarius Commission.
- Doyle, E. 2006. Veterinary drug residues in processed meats-potential health risk. A review of the scientific literature. Food Research Institute, University of Wisconsin-Madison, USA.
- EFSA. 2007. Annual EU pesticide residues monitoring report. The European Food Safety Authority.
- FSIS. 2000. Blue Book. 2000. FSIS National Residue Program. Food Safety and Institution Service, USDA.
- Huss, H., Ababouch, L., and Gram, L. 2004. Assurance of seafood quality. Danish Institute for Fisheries. Research Department of Seafood Research. Food and Agriculture Organization of the United States, Rome, Italy.
- Kroes, R., Muller, D., Lambe, J., Lowik, M., van Klaveren, J., Kleiner, J., Massey, R., Mayer, S., Urieta, I., Verger, P. and Visconti, A. 2002. Assessment of intake from the diet. Food and Chemical Toxicology. 40, 327-85.
- Kelly, L. 2005. Risk analysis framework. Risk analysis, application to chemicals. Food Standards Australia and New Zealand.
- FNRI. 2003. Food consumption survey. Food and Nutrition Research Institute. Philippines.
- Health Canada. 2003. Setting standards for maximum residue limits (MRLs) of veterinary drugs used in food-producing animals. Ontario, Canada.
- NMIS. 2009. Veterinary Drug Residue Monitoring Program. Veterinary Drug Residue Section, Laboratory Services Division, National Meat Inspection Service, Department of Agriculture, Philippines.
- Pocas, M. and Hogg, T. 2007. Exposure assessment of chemicals from packaging materials.
- RITM. 2009. The report on antimicrobial resistant surveillance program. Research Institute for Tropical Medicine. Philippines.
- Steve, J.C. 2008. Background paper on dietary exposure assessment. Prepared for the WHO Expert Meeting on Toxicological and Health Aspects of Melamine and Cyanuric Acid. Geneva.
- Sumner, J., Ross, T., and Ababouch, L. 2004. Application of risk assessment in the fish industry. Fisheries technical paper, 442. Food and Agriculture Organization of the United Nations. Rome, Italy.
- Titus, S. 2007. A NOEL model developed for quantitative microbial risk assessment in the pork food chain. A dissertation of Doctor of Philosophy at Massey University, Palmerston North, New Zealand.
- Venn, V., Loinda, R.B., Billy P.D., and RioJohn T.D. 2010. Dietary exposure assessment of antibiotic residues in pork consumed in the Philippines. MVM thesis. 199.
- WHO. 1999. Joint FAO/NACA/WHO study group on food safety issues associated with products from aquaculture. WHO Technical Report Series No. 883. World Health Organization, Geneva, Switzerland.
- Dong, Z., Li N., Wang Z.T., Zho Y.F., Wu Y.N., and Yan W.X. 2009. Assessment on dietary melanine exposure from tainted infant formula. 100-103.